

## CHECK LIST FOR PATIENTS WITH SEPTIC SHOCK

1. DIAGNOSIS      ↑      most likely cause \_\_\_\_\_
2. IV ACCESS      2 large bore cannulae ↑      where \_\_\_\_\_  
I/O access      ↑      where \_\_\_\_\_
3. MONITORING      ECG ↑ PR ↑ BP ↑ RR ↑ GCS/AVPU ↑  
O2 sats ↑ Pupils ↑ Urine catheter ↑
4. BLOOD TESTS      FBC ↑ U+E ↑ LFT ↑ Clotting ↑ Glucose ↑  
Ca/Mg/Po4 ↑ ABG ↑ Lactate ↑ CRP ↑ G+S ↑
5. CULTURES      Blood ↑ Urine ↑ ETT ↑ Stool ↑  
NPA ↑ PCR ↑
6. FLUIDS      Bolus #1      10 / 20 ml/kg    time given \_\_\_\_\_  
Bolus #2      10 / 20 ml/kg    time given \_\_\_\_\_  
Bolus #3      10 / 20 ml/kg    time given \_\_\_\_\_  
Bolus #4      10/20 ml/kg    time given \_\_\_\_\_  
Total fluid given \_\_\_\_\_ ml/kg
7. CONTACTED PICU      time phoned \_\_\_\_\_
8. INTUBATED      Etomidate/ Ketamine      Suxamethonium ↑  
Rocuronium ↑  
Other drugs used \_\_\_\_\_  
Tube size \_\_\_\_\_      Checked on CXR ↑
9. CENTRAL LINE      yes/no      if yes where \_\_\_\_\_
10. ARTERIAL LINE      yes/no      if yes where \_\_\_\_\_
11. SEDATION      Morphine ↑      Midazolam ↑  
Other \_\_\_\_\_

